

LSU Health Science Center

LSU Health Digital Scholar

Medical Research Day

2022 Medical Research Day Posters

Oct 13th, 12:00 AM

Risk of Malignancy Following Radiofrequency Ablation of Thyroid Nodules

Peter P. Issa

LSU Health Sciences Center- New Orleans

M Omar

Tulane University

C P. Issa

LSU Health Sciences Center- New Orleans

Y Buti

Tulane University

M Hussein

Tulane University

See next page for additional authors

Follow this and additional works at: <https://digitalscholar.lsuhscc.edu/sommrd>



Part of the [Surgery Commons](#)

Recommended Citation

Issa, Peter P.; Omar, M; Issa, C P.; Buti, Y; Hussein, M; Aboueisha, M; Abdelhady, A; Shama, M; Lee, G S.; Toraih, E; and Kandil, Emad, "Risk of Malignancy Following Radiofrequency Ablation of Thyroid Nodules" (2022). *Medical Research Day*. 36.

<https://digitalscholar.lsuhscc.edu/sommrd/2022MRD/Posters/36>

This Event is brought to you for free and open access by the School of Medicine at LSU Health Digital Scholar. It has been accepted for inclusion in Medical Research Day by an authorized administrator of LSU Health Digital Scholar. For more information, please contact aolini@lsuhscc.edu.

Presenter Information

Peter P. Issa, M Omar, C P. Issa, Y Buti, M Hussein, M Aboueisha, A Abdelhady, M Shama, G S. Lee, E Toraih, and Emad Kandil

Introduction

- Thyroid nodules can be classified as benign, malignant, or indeterminate, the latter of which make up 10–30% of nodules.
- Radiofrequency ablation (RFA) has become an attractive and promising therapy for the treatment of benign thyroid nodules.
- Though an attractive treatment modality, the current ATA guidelines recommend RFA only for the management of benign thyroid nodule, necessitating surgical management for the treatment of both indeterminate nodules (10–30% of all nodules) and malignant nodules (5% of all nodules).
- International literature, however, suggests RFA to be promising beyond their typical treatment of benign nodules, including papillary thyroid carcinomas.

Research Question

Can RFA safely and effectively treat indeterminate thyroid nodules?

Methods

- Our protocol consisted of preoperative FNA as well as comprehensive neck ultrasounds, treatment by RFA, and subsequent follow-up appointments at the 1, 3, 6, and 12 month marks. All operations were performed by fellowship-trained endocrine surgeons.
- VRR was calculated as $[(V_0 - V_1) / V_0] \times 100$, where V_0 signifies the initial nodule volume and V_1 the post-ablation nodule volume. VRR was utilized to determine operative success, defined as a VRR of $>50\%$ at final follow-up. Nodular regrowth was described as a nodule that was assessed by ultrasound to be of greater volume postoperatively than pre-operatively.

Results

Table 2: Demographic characteristics of included patients

Characteristics	Levels	Total	Bethesda III & IV nodules	Bethesda II nodules	p-value
Number		178	53	125	
Demographic data					
Age	Median (IQR)	64 (53-69)	63 (56-68)	65 (52.5-70)	0.49
Gender	Female	128 (71.9)	30 (56.6)	98 (78.4)	0.003
	Male	50 (28.1)	23 (43.4)	27 (21.6)	
Race	African American	100 (56.2)	28 (52.8)	72 (57.6)	0.82
	White	71 (39.9)	23 (43.4)	48 (38.4)	
BMI	Median (IQR)	30 (26.7-34.3)	30.5 (27.5-34.5)	29.8 (26.4-34.3)	0.38
Baseline sonographic features					
Nodule maximum diameter	Median (IQR)	2.4 (1.5-4)	2.5 (1.5-4.3)	2.3 (1.4-3.7)	0.47
Baseline Volume	Median (IQR)	1.8 (1.1-2.9)	1.9 (1.1-3.2)	1.7 (1.1-2.7)	0.45
	Solid	5 (2.8)	3 (5.7)	2 (1.6)	
Composition	Cystic	157 (88.2)	46 (86.8)	111 (88.8)	0.31
	Mixed	16 (9)	4 (7.5)	12 (9.6)	
	Hypoechoic	33 (21.3)	11 (23.4)	22 (20.4)	
Echogenicity	Isoechoic	121 (78.1)	36 (76.6)	85 (78.7)	0.74
	Hyperechoic	1 (0.6)	0 (0)	1 (0.9)	
	Grade 0	11 (7.9)	1 (2.3)	10 (10.3)	
Vascularity	Grade 1	46 (32.9)	11 (25.6)	35 (36.1)	0.054
	Grade 2	53 (37.9)	23 (53.5)	30 (30.9)	
	Grade 3	30 (21.4)	8 (18.6)	22 (22.7)	
	Soft	7 (5.3)	4 (10)	3 (3.2)	
Elastography	Mixed	96 (72.2)	27 (67.5)	69 (74.2)	0.27
	Stiff	30 (22.6)	9 (22.5)	21 (22.6)	
Calcifications	No Calcifications	91 (58.7)	29 (61.7)	62 (57.4)	0.25
	Microcalcifications	51 (32.9)	12 (25.5)	39 (36.1)	
	Macrocalcifications	13 (8.4)	6 (12.8)	7 (6.5)	
Laboratory data					
Baseline TSH uIU/mL	Median (IQR)	1.3 (0.7-1.9)	1.3 (0.8-1.9)	1.3 (0.7-2)	0.67
Post procedural TSH uIU/mL	Median (IQR)	1.2 (0.7-1.8)	1.1 (0.7-1.8)	1.2 (0.8-1.8)	0.58




Figure 1: Benign and indeterminate thyroid nodules treated by RFA (A) success rates and (B) regrowth rates. Success was a volume reduction rate of $\geq 50\%$. Nodular regrowth was defined as a nodule which was assessed by ultrasound to be of greater volume postoperatively than pre-operatively.

Findings

- 178 patients with thyroid nodules diagnosed as benign (Bethesda II) or indeterminate (Bethesda III/IV) by preoperative cytopathological analysis were included.
- Though patients with benign nodules tended to be female, patients were similar with respect to age, race, BMI, baseline sonographic features, and pre-operative biochemical parameters.
- Patients in the benign and indeterminate cohorts had similar thyroid nodule volume reduction rates at 65.60% and 64.20%, respectively ($p=0.68$).
- The two groups had similar nodular regrowth rates, at 11.2% for benign nodules and 9.40% for indeterminate nodules ($p=0.72$).
- A total of 3 cases of transient dysphonia were reported.

Discussion

- RFA of indeterminate thyroid nodules was comparable to that of benign thyroid nodules in all parameters of interest, including volume reduction rate.
- To our best knowledge, our work is the first North American analysis comparing benign and indeterminate thyroid nodules and suggests RFA to be a promising modality for the management of indeterminate thyroid nodules.
- Future larger studies are warranted to further elucidate the difference, if any, in managing different nodule Bethesda classes.

References

- Dean, D.S.; Gharib, H. Fine-Needle Aspiration Biopsy of the Thyroid Gland. Endotext Internet 2015.
- Haugen, B.R.; Alexander, E.K.; Bible, K.C.; Doherty, G.M.; Mandel, S.J.; Nikiforov, Y.E.; Pacini, F.; Randolph, G.W.; Sawka, A.M.; Schlumberger, M. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. Thyroid 2016, 26, 1–133.
- Ha, S.M.; Sung, J.Y.; Baek, J.H.; Na, D.G.; Kim, J.; Yoo, H.; Lee, D.; Whan Choi, D. Radiofrequency Ablation of Small Follicular Neoplasms: Initial Clinical Outcomes. Int. J. Hyperthermia 2017, 33, 931–937.