

LSU Health Science Center

## LSU Health Digital Scholar

---

Medical Research Day

2022 Medical Research Day Posters

---

Oct 13th, 12:00 AM

# The Role of TRAF3IP2 in the Progression of Alcoholic Cardiomyopathy

Nicholas R. Harris

*LSU Health Sciences Center- New Orleans*

Joshua M. Edavettal

*LSU Health Sciences Center- New Orleans, jedav1@lsuhsc.edu*

Jason D. Gardner

*LSU Health Sciences Center- New Orleans, jgardn@lsuhsc.edu*

Follow this and additional works at: <https://digitalscholar.lsuhscc.edu/sommrd>



Part of the [Medical Physiology Commons](#)

---

### Recommended Citation

Harris, Nicholas R.; Edavettal, Joshua M.; and Gardner, Jason D., "The Role of TRAF3IP2 in the Progression of Alcoholic Cardiomyopathy" (2022). *Medical Research Day*. 95.

<https://digitalscholar.lsuhscc.edu/sommrd/2022MRD/Posters/95>


This Event is brought to you for free and open access by the School of Medicine at LSU Health Digital Scholar. It has been accepted for inclusion in Medical Research Day by an authorized administrator of LSU Health Digital Scholar. For more information, please contact [aolini@lsuhsc.edu](mailto:aolini@lsuhsc.edu).

## Introduction


**Alcoholic Cardiomyopathy (ACM)** manifests in humans after excessive alcohol consumption and is characterized by ventricular dilation and cardiac function impairment. Previous studies have identified deterioration of mitochondrial homeostasis, increased oxidative stress, and inflammation as mechanisms of ACM development.<sup>1</sup> However, little is known of the molecular mechanism of ACM. Here, we focus on the role of **TRAF3IP2**, a proinflammatory cytoplasmic adapter protein, in the pathogenesis of ACM. Previous work suggests TRAF3IP2 is a master regulator of inflammation; thus, being a potential therapeutic target.

## Chronic-Binge ETOH Model

In this study, we used a mouse model of chronic plus binge alcohol feeding described by the NIAAA<sup>2</sup>. After 5 days of acclimation to the liquid diet, mice are fed *ad libitum* 5% ethanol (EtOH) liquid diet (Lieber-DeCarli) or isocaloric control liquid diet for either 10 days or 20 days. At days 10 and 20, mice received an oral binge dose of EtOH (5 g/kg body wt), or isocaloric maltose dextrin solution (9 g/kg body wt), via oral gavage. A graphical timeline<sup>3</sup> is shown below:



## Echocardiography



## qPCR






Table 1								
P-values	COL1A1	COL3A1	IL-1 $\beta$	IL-6	LARP6	LOX	TGF- $\beta$	TRAF3IP2
10d	0.06	0.08	0.88	0.78	<b>0.04*</b>	0.24	0.91	0.05
20d	0.82	0.25	0.36	0.26	0.89	0.42	0.90	0.26

**Figures 2 and 3:** RNA isolated from the left ventricle was analyzed via qPCR. Gene expression was normalized using 18S rRNA as a reference gene. Student's *t*-test was used to assess difference in gene expression between ethanol- and control-diet cohorts. A *p*-value  $< 0.05$  was considered significant and results are presented as mean  $\pm$  Std. *P*-values are shown in **Table 1**.

## Invasive Hemodynamics

After either 10 days or 20 days, left ventricular function was assessed via catheterization with a pressure-volume conductance catheter. End systolic pressure volume relationship (**ESPVR**) and preload recruitable stroke work (**PRSW**) are indices of contractility.



**Figure 4:** PV Loop with calculated ESPVR, 20d Control Mouse. **Figure 5:** Stroke work vs End-diastolic volume with calculated PRSW, 20d Control Mouse. **Figure 6:** PV Loop with calculated ESPVR, 20d ETOH mouse. **Figure 7:** Stroke work vs End-diastolic volume with calculated PRSW, 20d ETOH mouse.

## Results

- Ethanol mice at 10 days had an increased Ejection Fraction.
- LARP6 was increased following 10 days of ethanol exposure and binge.
- There were no significant differences in qPCR and echocardiography measurements between control and ethanol mice at 20 days.

## Discussion

Results from this study showed little differences in gross and molecular cardiac structure and function. Further studies are needed to elucidate the role of TRAF3IP2 in the development of ACM.

## References

- Matyas et al. "Chronic plus Binge Ethanol Feeding Induces Myocardial Oxidative Stress, Mitochondrial and Cardiovascular Dysfunction, and Steatosis." *American Journal of Physiology-Heart and Circulatory Physiology* 310, no. 11 (June 2016): H1658-70.
- Bertola, et al. "Mouse Model of Chronic and Binge Ethanol Feeding (the NIAAA Model)." *Nature Protocols* 8, no. 3 (March 2013): 627-37.
- Created with Biorender.