

LSU Health Science Center

LSU Health Digital Scholar

Medical Research Day

2022 Medical Research Day Posters

Oct 13th, 12:00 AM

Use of Viscoelastic Testing in Trauma Patients

Nicolas Chanes

LSU Health Sciences Center- New Orleans

Shyam Murali

University of Pennsylvania

Eric Winter

University of Pennsylvania

Madhu Subramanian

University of Pennsylvania

Allyson Hynes

University of Pennsylvania

See next page for additional authors

Follow this and additional works at: <https://digitalscholar.lsuhscc.edu/sommrd>



Part of the [Surgery Commons](#)

Recommended Citation

Chanes, Nicolas; Murali, Shyam; Winter, Eric; Subramanian, Madhu; Hynes, Allyson; Seamon, Mark; Cannon, Jeremy; and Smith, Alison, "Use of Viscoelastic Testing in Trauma Patients" (2022). *Medical Research Day*. 91.

<https://digitalscholar.lsuhscc.edu/sommrd/2022MRD/Posters/91>

This Event is brought to you for free and open access by the School of Medicine at LSU Health Digital Scholar. It has been accepted for inclusion in Medical Research Day by an authorized administrator of LSU Health Digital Scholar. For more information, please contact aolini@lsuhscc.edu.

Presenter Information

Nicolas Chanes, Shyam Murali, Eric Winter, Madhu Subramanian, Allyson Hynes, Mark Seamon, Jeremy Cannon, and Alison Smith

Introduction

- Approximately 25% of trauma patients develop some form of coagulopathy that impacts their clinical course (1).
- Trauma-induced coagulopathy (TIC) carries significant risk, with TIC+ patients experiencing up to three times the mortality rates of TIC- patients (2).
- Despite awareness of this increased mortality and improvements in damage control resuscitation, the etiology of TIC has not been fully elucidated and is likely multifactorial (3).
- Currently, the management of TIC relies on standardized treatment protocols that do not address the nature of the hemostatic deficits, but rather provide blood products in proportions that approach the composition of whole blood (4).
- Thromboelastography (TEG) and rotational thromboelastometry (ROTEM), which provide a point-of-care assessment of global clot formation and dissolution, have become an integral part of the management of trauma patients as they can be used in real time to guide the transfusion of blood products.
- Recent investigations have suggested that the use of TEG/ROTEM testing may be of value in the early management of trauma patients; however, there is a need for determining the specific TEG/ROTEM criteria and thresholds that guide hemostatic resuscitation and predict mortality (5).

Objectives

- Conduct a retrospective review of a pre-existing database containing adult trauma patients who required massive transfusion protocol and had viscoelastic testing (TEG or ROTEM) in order to develop a TEG score that may be used to better guide future practice.
- Use the findings we obtain from this study to determine the specific TEG/ROTEM criteria and thresholds that guide hemostatic resuscitation and predict mortality.

Methods

- This was a secondary analysis of a multicenter retrospective review, which included adult patients receiving massive transfusion protocol at 7 level 1 trauma centers (2012-2018).
- TIC was defined using a "TEG score" with one point assigned for abnormalities in each of the following:
 - Clot time (>8.9 min)
 - Alpha angle (<65 degrees)
 - Clot strength (<55 mm).
- TIC+ patients (TEG score 1-3) were compared to TIC- patients (TEG score 0).
- Multiple regression was used to control confounding variables while evaluating the association between abnormal TEG values and the endpoints of in-hospital mortality (primary outcome) and 24-hour blood product transfusion (secondary outcome).

ROTEM and TEG Explained






Figure 1. A typical ROTEM trace (top) and TEG trace (bottom). Adapted from Sinclair, et. al (6).

Results



Results (con't)

- Of 1,145 total patients, 442 (38.6%) were TIC+.
- On adjusted analysis, abnormal clot time significantly predicted 24-hour platelet ($p<0.001$) and cryoprecipitate ($p=0.046$) transfusion requirements.
- Abnormal alpha-angle similarly predicted 24-hour platelet ($p<0.001$) and cryoprecipitate ($p<0.001$) transfusion.
- Abnormal clot strength predicted in-hospital mortality ($p=0.015$) and 24-hour fresh frozen plasma transfusion ($p=0.016$).
- Increasing TEG score significantly predicted mortality ($p<0.001$), 24-hour packed red blood cell ($p=0.028$), and 24-hour platelet transfusion ($p<0.001$).

Conclusions and Future Directions

Conclusions

- Abnormal TEG values are independently associated with greater transfusion requirements and increased mortality.
- A TEG score based on the number of abnormal parameters reliably identifies TIC and an increased score is associated with greater TIC severity.

Future Directions

- Further research is needed to validate the TEG score in a prospective manner.

References

- Branco BC, David JS, et al. Thromboelastogram evaluation of the impact of hypercoagulability in trauma patients. *Shock*. 2014;41(3):200-7.
- Frith D, Allard S, et al. Definition and drivers of acute traumatic coagulopathy: clinical and experimental investigations. *J Thromb Haemost*. 2010;8(9):1919-25.
- Madurska MJ, Morrison JJ, et al. Fibrinolysis in trauma: a review. *Eur J Trauma Emerg Surg*. 2018;44(1):35-44.
- Baksaas-Aasen K, Maegele M, et al. Viscoelastic haemostatic assay augmented protocols for major trauma haemorrhage (ITACTIC): a randomized, controlled trial. *Intensive Care Med*. 2021;47(1):49-59.
- Veigas PV, da Luz LT, et al. A systematic review on the rotational thromboelastometry (ROTEM®) values for the diagnosis of coagulopathy, prediction and guidance of blood transfusion and prediction of mortality in trauma patients. *Scand J Trauma Resusc Emerg Med*. 2016;24(1):114.
- Anderson, L., Quasim, I., Steven, M., Moise, S. F., Shelley, B., Schraag, S., & Sinclair, A. (2014). Interoperator and intraoperator variability of whole blood coagulation assays: a comparison of thromboelastography and rotational thromboelastometry. *Journal of cardiothoracic and vascular anesthesia*, 28(6), 1550–1557. <https://doi.org/10.1053/j.jvca.2014.05.023>