

LSU Health Science Center

LSU Health Digital Scholar

Medical Research Day

2021 Medical Research Day Posters

Oct 19th, 12:00 AM

Demographics of Periprosthetic Joint Infection Rates

Katelynn Donnelly

LSU Health Sciences Center- New Orleans

Abigail Erwin

LSU Health Sciences Center- New Orleans, aerwi1@lsuhsc.edu

Andrew Chapple

LSU Health Sciences Center- New Orleans, achapp@lsuhsc.edu

Peter Krause

LSU Health Sciences Center- New Orleans, PKraus@lsuhsc.edu

Vinod Dasa

LSU Health Sciences Center- New Orleans, vdasa@lsuhsc.edu

Follow this and additional works at: <https://digitalscholar.lsuhscc.edu/sommrd>



Part of the [Medical Biomathematics and Biometrics Commons](#)

Recommended Citation

Donnelly, Katelynn; Erwin, Abigail; Chapple, Andrew; Krause, Peter; and Dasa, Vinod, "Demographics of Periprosthetic Joint Infection Rates" (2021). *Medical Research Day*. 27.

<https://digitalscholar.lsuhscc.edu/sommrd/2021MRD/Posters/27>

This Event is brought to you for free and open access by the School of Medicine at LSU Health Digital Scholar. It has been accepted for inclusion in Medical Research Day by an authorized administrator of LSU Health Digital Scholar. For more information, please contact aolini@lsuhsc.edu.

Demographics of Periprosthetic Joint Infection Rates

Katelynn Donnelly, MS, Abigail Erwin, BS, Andrew Chapple, PhD, Peter C. Krause, MD, Vinod Dasa, MD



1. LSUHSC, Biostats Program, School of Public Health 2. LSUHSC, Department of Orthopaedics

Introduction

Risk factors for periprosthetic joint infections (PJIs) following total joint arthroplasties (TJAs) include patient demographics and comorbidities such as age, gender, obesity, rheumatoid arthritis, and diabetes.¹ However, there is insufficient research analyzing the number of PJIs in certain racial subsets of the population despite the presence of disparities due to race that still exists in orthopedics today.² The purpose of this study is to investigate if PJI rates differ by race, as well as other patient demographics. This may uncover valuable information on how race relates to PJIs which is frequently unaccounted for in other studies. This study aims to analyze the current data to highlight possible disparities in PJI prevalence due to racial background.

Results




Figure 1: Out of the total number of patients included in the study (21,735), a total of 746 patients developed a PJI following a TJA. Out of the 746 patients, 82% were white, 16% were black, and 2% were of another race.




Figure 2: Out of the total number of patients included in the study (21,735), a total of 746 patients developed a PJI following a TJA. Out of the 746 patients, 9% had private insurance and 91% had public insurance.

Results Continued

Variable	OR (CI)	P-value
Black v White	0.72 (0.58-0.88)	0.002
Other Race v White	0.8 (0.45-1.42)	0.444
Hispanic Ethnicity	1.21 (0.86-1.7)	0.277
Alcohol Use	1.83 (1.3-2.57)	0.001
Smoking	1.19 (1.02-1.38)	0.028
Private v Public Insurance	0.59 (0.45-0.79)	<.001
Age (Continuous)	0.97 (0.96-0.98)	<.001
CCI (Continuous)	1.21 (1.16-1.25)	<.001
Year (Continuous)	1.52 (1.45-1.6)	<.001




Figure 4: This figure displays a multivariable logistic regression analysis for patient's with PJI development in one year.

Methods

This retrospective cohort extracted patient data from Reachnet for patients with procedure codes for THA, TKA, and TSA, as well as diagnostic codes for PJIs to determine rates of PJIs by patient demographics. Patient demographic data included race, alcohol use, smoking, insurance, BMI > 30, age ≥ 70, procedure year, and Charlson Comorbidity Index. Patients without a follow-up within one year, missing BMI, or non-public/private insurance were excluded. All in all, this resulted in 21,735 patients total. Percents, counts, multivariable logistic regressions were used to summarize the data collected.

Percentage of PJI Development in Patients with Comorbid Conditions




Figure 3: This figure displays frequency of PJI development in patients who used alcohol, smoked, had a BMI >30, or were over the age of 70. In those who used alcohol, about 7% developed a PJI post TJA. In those who were smokers, had a BMI >30, or were over the age of 70 about 3% developed a PJI.

Conclusions

- From our specific cohort, black patients had a significantly decreased risk of PJI within 1 year (Odds ratio = .72, 95% CI=.58-.88) compared to white patients.
- Other demographics such as alcohol use, smoking, and increased Charlson Comorbidity Index contributed to considerably higher rates.
- Additionally, rates of PJIs have significantly increased in recent years.
- Increased age and private insurance were associated with decreased rates of PJI.

References

- 1.Kong L, Cao J, Zhang Y, Ding W, Shen Y. Risk factors for periprosthetic joint infection following primary total hip or knee arthroplasty: a meta-analysis. *Int Wound J.* 2017;14(3):529-536. doi:10.1111/iwj.12640
- 2.Singh JA, Lu X, Rosenthal GE, Ibrahim S, Cram P. Racial disparities in knee and hip total joint arthroplasty: an 18-year analysis of national Medicare data. *Ann Rheum Dis.* 2014;73(12):2107-2115. doi:10.1136/annrheumdis-2013-203494